

Species

To Cite:

Li Y, Elie N, Tao X, Zhang H. New Record of *Rhipidothripiella* Bagnall, 1932 (Thysanoptera, Aeolothripidae) from Ethiopia, the North East of Africa. *Species* 2024; 25: e44s1723
doi: <https://doi.org/10.54905/disssi.v25i76.e44s1723>

Author Affiliation:

¹Plant Protection College, Yunnan Agricultural University, Kunming, Yunnan, China

²Haramaya University, ACE Climate SABC, P.o. Box:138 Harar, Haramaya, Ethiopia

³University of Rwanda, CST & CoEB, KN 67 Street Nyarugenge, Kigali, Rwanda

Corresponding Author

Plant Protection College, Yunnan Agricultural University, Kunming, Yunnan, China
Email: hongruizh@126.com

ORCID List

| | |
|------------------|---------------------|
| Yajin Li | 0000-0003-3815-3021 |
| Ntirenganya Elie | 0000-0002-4603-5693 |
| Xiran Tao | 0009-0006-6390-7613 |
| Hongrui Zhang | 0000-0002-0089-1099 |

Peer-Review History

Received: 02 July 2024
Reviewed & Revised: 06/July/2024 to 03/October/2024
Accepted: 07 October 2024
Published: 10 October 2024

Peer-Review Model

External peer-review was done through double-blind method.

Species

pISSN 2319-5746; eISSN 2319-5754



© The Author(s) 2024. Open Access. This article is licensed under a [Creative Commons Attribution License 4.0 \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.



New Record of *Rhipidothripiella* Bagnall, 1932 (Thysanoptera, Aeolothripidae) from Ethiopia, the North East of Africa

Yajin Li¹, Ntirenganya Elie^{2,3}, Xiran Tao¹, Hongrui Zhang^{1*}

ABSTRACT

The species of *Rhipidothripiella turneri* (Moulton, 1930) is newly recorded from Ethiopia. This species is detail described, a list of 7 genera and 25 Aeolothripidae species in Africa is provided.

Keywords: Metanotal sculpture, sensorium, *Rhipidothripiella*, grasses

1. INTRODUCTION

More than 200 species of Aeolothripidae are recognized in 23 genera around the world, and more than 50% of them are included in the Holarctic genus *Aeolothrips* Haliday and Australian genus *Desmothrips* Hood. Seven monotypic genera were distributed in Africa (*Corynothripoides* Bagnall, *Euceratothrips* Hood, *Rhipidothripiella* Bagnall), Australia (*Andrewarthaia* Mound, *Erythridothrips* Mound & Marullo, *Lamprothrips* Moulton) and Asia (*Indothrips* Bhatti). Mound and Marullo, (1998) discussed the diversity of biologies amongst Aeolothripidae and provided a key of 36 species in 12 genera in Australia. Masumoto and Okajima, (2019) studied 3 genera and 10 species of Aeolothripidae in Japan. This family in Africa is poorly studied, most species were recorded in the last century and with few detailed informations.

Mound, (1968) revised Bagnall's collection and redefined several genera and a few species. Most Aeolothripidae species are distinguished by the linear sensoria on antennal segments III-IV and curved around the apex of the segment, still, members of the *Rhipidothrips* Uzel genus are marked by the sensoria retained only the distal curved portion, while some species of *Cycadothrips* Mound, *Dactuliothrips* Moulton and *Orothrips* Moulton segments III-IV posse two sensoria (Marullo and Mound, 1993). Metanotum sculpture with little difference among different genera in this family (from Laurence Mound's personal communications, 2024), the relationships of these genera are unknown. To provide enough information for further study about the Aeolothripidae species, the species of *Rhipidothripiella turneri* collected from *Calamagrostis epigejos* [Grass: Poaceae] is described.

2. MATERIALS AND METHODS

The specimens were collected in Haramaya University (9° 25' 23" N, 42°02' 14" E), Haramaya district, this area is located in the eastern part of Ethiopia, it is characterized by a sub-humid type of climate. The annual mean temperature is 9.9-24.18°C, whereas the annual mean rainfall ranges between 600-1260 mm. The short and main rain seasons extend from March to May and July to September, respectively (Sorecha, 2017). Thrips collection and slide preparation methods followed (Zhang et al., 2006). Specimens were observed with an Olympus BX53 System Microscope. Figures were made using an Olympus DP74 CCD with cellSens imaging software. Slide-mounted specimens are deposited at Yunnan Agricultural University, Kunming.

3. RESULTS AND DISCUSSION

Rhipidothripiella turneri (Moulton, 1930)

Female. Macropterous. Body uniformly brown (Figure 1); antennae brown, III- IV yellow with apex of IV light brown (Figure 2G); fore wing with two dark cross bands alternating with two pale areas, the pale areas at base and sub-apex, clavus pale (Figure 2F); all legs dark brown; prominent body setae brown. Head longer than wide, slightly produced in front of eyes, with transverse close striae, postocular setae in two rows but irregularly arranged (Figure 2A); eyes extending posteriorly on ventral side; maxillary palp 3-segmented, II subdivided into five parts, the structures between the fourth and the fifth is distinct (Figure 2C). Antennal segments III-IV with sensoria vermicular, recurved and crossing around apex on one side and turn straight towards the base on the other side, sometimes sensorium on III with an interruption medially (Figure 2H), segment III slightly more extended than IV, VI-VII almost subequal (Figure 2G).



Figure 1 *Rhipidothripiella turneri*. Body of Female.

Pronotum is wider than long, smooth, with about 30 discal setae and three pairs of posteromarginal setae, one pair of posteroangular setae slightly longer than discal setae (Figure 2A). Mesonotum sculptured with closely transverse striae, with one pair of setae medially, anteromedian CPS absent (Figure 2D). Metascutum with weak reticulations medially, poster and lateral part sculptured with longitudinal striae, one pair of small setae near posterior, CPS present (Figure 2D). Spinula absent on both meso- and metasternum (Figure 2E). Fore tarsus with two stout setae (Figure 2I).




Figure 2 *Rhipidothripiella turneri*. A. Head and pronotum. B. Abdominal sternites V–VII. C. Ventral of head and pronotum. D. Mesonotum and metanotum. E. Mesosternum and metasternum. F. Fore wing. G. Antennae. H. Antennal segments III–IV. I. Fore tibiae.

Fore wing first and second vein with setal rows regular and complete, clavus with six anteromarginal setae (Figure 2F). Abdominal sternites I-VII with two pairs of posteromarginal setae, the setae at the posterior margin except segment I in front of margin; sternites II-IV each with two pairs of discal setae laterally, V-VII with four pairs of discal setae laterally; VII with two pairs of accessory setae between S1 and S2 (Figure 2B).

Materials examined

Ethiopia, Harar, Haramaya University Road; 9° 25' 23" N and 42° 02' 14" E; alt.2057m; 11.X.2022; Ntirenganya Elie leg.; 7 ♀, collected from *Calamagrostis epigejos* [Poaceae].

Discussion

Moulton, (1930) described *Rhipidothrips turneri* species from South Africa with only a single male specimen. Bagnall, (1932) built *Rhipidothripiella* genus and erected *turneri* as a type species. This monotypic genus was related to *Allelothrips* Bagnall than *Rhipidothrips* Moulton with pronotum has no posteroangular setae, but the fore wings were not as Bagnall stated, and the metanotal sculpture is quite different from that of *Allelothrips* (Bagnall, 1932; Mound, 1968). The relationship of this genus is not clear, and more collections and profound studies are needed. This species is newly recorded in Ethiopia. The specimens were collected from all parts of *Calamagrostis epigejos* plants, and in association with leaf mites, which led to the predatory feeding habits suspect.

4. CONCLUSION

A list of 7 genera and 25 Aeolothripidae species in Africa are provided (Table 1), including *Rhipidothripiella turneri* Moulton, (1930) was newly recorded in Ethiopia.

Table 1 Aeolothripidae species from Africa

| Genera | Species | Distribution | References |
|--------------------------------------|------------------------------------|----------------------------------|-------------------------------------|
| <i>Aeolothrips</i> Haliday, 1836 | <i>arnebiae</i> Priesner, 1948 | Sudan | Priesner, 1948 |
| | <i>brevicornis</i> Bagnall, 1915 | South Africa | Moulton, 1930 Mound, 1968 |
| | <i>collaris</i> Priesner, 1919 | Tunisia | Kort et al., 2020 Priesner, 1938 |
| | <i>deserticola</i> Priesner, 1929 | Egypt, Iran Israel Morocco | Priesner, 1948 |
| | <i>fasciatus</i> (Linnaeus, 1758) | South Africa | Moulton 1936 |
| | <i>intermedius</i> Bagnall, 1934 | Tunisia | Kort et al., 2020 |
| | <i>scabiosatibia</i> Moulton, 1930 | South Africa | Moulton, 1930 |
| | <i>tenuicornis</i> Bagnall, 1926 | Tangier | Mound, 1968 |
| | <i>variicornis</i> Bagnall, 1932 | Tanganyika | Bagnall, 1932 Mound, 1968 |
| | <i>wittmeri</i> Priesner, 1935 | Egypt | Priesner, 1948 |
| <i>Allelothrips</i> Bagnall, 1932 | <i>brunneus</i> (Faure, 1941) | South Africa | Faure, 1941 |
| | <i>cameroni</i> (Bagnall, 1932) | Sudan | Bagnall, 1932 |
| | <i>cincticornis</i> Bagnall, 1932 | Sudan | Bagnall, 1932 |

| | | | |
|--|---|--|---|
| | <i>talithae</i> (Faure, 1941) | South Africa | Faure, 1941 |
| | <i>tenuicornis</i> (Bagnall, 1932) | Sudan | Mound, 1968 |
| | <i>vilardeboi</i> (Bournier & Bournier, 1979) | South Africa | Mound and Marullo, 1993 |
| <i>Audiothrips</i> Moulton, 1930 | <i>perplexus</i> Moulton, 1930 | South Africa | Mound and Marullo, 1993 |
| | <i>senckenbergiana</i> Mound & Marullo, 1993 | South Africa | Mound and Marullo, 1993 |
| <i>Corynothripoides</i> Bagnall, 1926 | <i>marginipennis</i> Bagnall, 1926 | Sierra Leone, Zaire, Congo, Rwanda, Uganda | Mound, 1968 Bagnall, 1926 Mound and Reynaud, 2005 |
| <i>Franklinothrips</i> Back, 1912 | <i>atlas</i> Hood, 1957 | Congo | Hood, 1957 |
| | <i>megalops</i> (Trybom, 1912) | South Africa | Bagnall, 1926 Moulton, 1930 Kort et al., 2020 |
| | <i>vespiformis</i> (Crawford, 1909) | South Africa | Bagnall 1926 Kort et al., 2020 |
| | <i>tenuicornis</i> Hood, 1915 | South Africa | Bagnall, 1926 |
| <i>Mymarothrips</i> Bagnall, 1928 | <i>ritchianus</i> Bagnall, 1928 | Tanzania | Bagnall, 1928 |
| <i>Rhipidothripiella</i> Bagnall, 1932 | <i>turneri</i> (Moulton, 1930) | South Africa Ethiopia | Bagnall, 1932 Present paper |

Acknowledgements

Thanks to Laurence Mound (CSIRO, Australia) for providing references and helping us to identify the species. We are also thanks to the Editor and the anonymous reviewers for the valuable suggestions.

Author contributions

Conceptualization: Hongrui Zhang; Investigation: Ntirenganya Elie; Methodology: Xiran Tao; Resources: Ntirenganya Elie; Supervision: Hongrui Zhang; Project administration: Yajin Li, Ntirenganya Elie; Software: Xiran Tao; Writing – original draft: Yajin Li; Writing – review and editing: Ntirenganya Elie, Hongrui Zhang. All authors reviewed the manuscript.

Ethical approval & declaration

In this article, as per the animal regulations followed in University of Rwanda, Kigali, Rwanda; Haramaya University, Ethiopia & Yunnan Agricultural University, Yunnan. China, the authors observed & recorded the species *Rhipidothripiella turneri* Moulton, (1930) newly from Ethiopia. Slide-mounted specimens are deposited at Yunnan Agricultural University, Kunming, China. The Animal ethical guidelines are followed in the study for species observation, identification & experimentation.

Conflicts of interests:

The authors declare that there are no conflicts of interests.

Funding:

The study was supported by Yunnan Fundamental Research Projects (No. 202301AU070120), Scientific Research Funds of Department of Education of Yunnan Province (No. 2023J0463), and National Natural Science Foundation of China (No. 32360129) and was the side work of the second author, who is in the PhD program under World Bank-Korea Trust Fund agreement (No. TF0A8639) for the PASET Regional Scholarship and Innovation Fund (No. RSIF/RA/013).

Data and materials availability

All data associated with this study are present in the paper.

REFERENCES

1. Bagnall RS. XXI—The family Franklinothripidae nov., with description of a new type of Thysanopteron. *Ann Mag Nat Hist* 1926; 17(97):168–173. doi: 10.1080/00222932608633386
2. Bagnall RS. Descriptions of some new genera and species of African Aeolothripoid Thysanoptera. *Ann Mag Nat Hist* 1932; 10(57):287–294. doi: 10.1080/00222933208673576
3. Bagnall RS. Preliminary description of *Mymarothrips ritchianus*, a new type of Thysanopteron. *Ann Mag Nat Hist* 1928; 10(1):304–307. doi: 10.1080/00222932808672783
4. Faure JC. Records and descriptions of South African Thysanoptera-II. *J Entomol Soc South Africa* 1941; 4(1):100–110.
5. Hood JD. Two new Thysanoptera from the Belgian Congo. *Proc R Entomol Soc London. Series B, Taxonomy* 1957; 26(9–10):172–174. doi: 10.1111/j.1365-3113.1957.tb00399.x
6. Kort IB, Lourdes Moraza M, Attia S, Mansour R, Kheder SB. Beneficial arthropods as potential biocontrol candidates of thrips (Thysanoptera: Thripidae) occurring in Tunisian citrus orchards. *Biologia* 2020; 75:2261–2270. doi: 10.2478/s11756-020-00487-x
7. Marullo R, Mound LA. Discontinuous distribution and systematic relationships of the genus *Orothrips* (Thysanoptera: Aeolothripidae) and related taxa in Mediterranean climates. *J N Y Entomol Soc* 1993; 101(4):561–566.
8. Masumoto M, Okajima S. Review of the Aeolothripidae (Thysanoptera) in Japan. *Zootaxa* 2019; 4564(2):301–326. doi: 10.11646/zootaxa.4564.2.1
9. Moulton D. LII —Thysanoptera from Africa. *Ann Mag Nat Hist* 1936; 17(101):493–509. doi: 10.1080/00222933608655149
10. Moulton D. Thysanoptera from Africa. *Ann Mag Nat Hist* 1930; 10(5):194–207.
11. Mound LA, Marullo R. Biology and identification of Aeolothripidae (Thysanoptera) in Australia. *Invertebr Taxon* 1998; 12(6):929–950. doi: 10.1071/it97014
12. Mound LA, Marullo R. The *Erythrothrips* complex of tropical Aeolothripidae (Thysanoptera) with a new Australian genus and a new South Africa. *Insect Syst Evol* 1993; 24(3):285–291. doi: 10.1163/187631293x00118
13. Mound LA, Reynaud P. *Franklinothrips*; a pantropical Thysanoptera genus of antmimicking obligate predators (Aeolothripidae). *Zootaxa* 2005; 864:1–16. doi: 10.11646/zootaxa.864.1.1
14. Mound LA. A review of R.S. Bagnall's Thysanoptera collections. *Bull Br Mus (Nat Hist). Entomol Suppl* 1968; 11:1–181.
15. Priesner H. Contributions towards a knowledge of the Thysanoptera of Egypt, XIV. *Bulletin de la Societe Royale entomologique d' Egypte* 1948; 32:317–341.
16. Priesner H. Thysanopteren aus dem Belgischen Congo (4. Beitrag). *Revue de Zoologie et de Botanique Africaines* 1938; 30(3):343–355.
17. Sorecha EM. Climate Projection Outlook in Lake Haramaya Watershed, Eastern Ethiopia. *Hydrol Current Res* 2017; 8:275. doi: 10.4172/2157-7587.1000275
18. Zhang HR, Okajima S, Mound LA. Collecting and slide preparation methods of thrips. *Chin Bull Entomol* 2006; 43(5):725–728.